

POULTRY

VACCINATION PROGRAM

POULTRY

VACCINATION PROGRAM



BROILERS VACCINATION PROGRAM

Age	Vaccine	Disease	Route
1-3 days	Hipraviar-B1/H120	ND & IB	Eye drop or spray or drinking water
10 days	Hipragumboro-CH80 or Hipragumboro-GM97	Gumboro	Drinking water
18 days	Hipragumboro-CH80 or Hipragumboro-GM97	Gumboro	Drinking water
21 days	Hipraviar –S or Hipraviar CLON	NCD	Eye drop or spray or drinking water

LAYERS VACCINATION PROGRAM

Age	Vaccine	Disease	Route
1-3 days	Hipraviar-B1/H120	ND & IB	Eye drop or spray or drinking water
10 days	Hipragumboro-CH80 or Hipragumboro-GM97	Gumboro	Eye drop or Drinking water
18 days	Hipragumboro-CH80 or Hipragumboro-GM97	Gumboro	Drinking water
21 days	Hipraviar –S or Hipraviar CLON	NCD	Eye drop or spray or drinking water
28 days	Hipraviar –S or Hipraviar CLON	NCD	Eye drop /Spray or drinking water
42 days	Hipraviar-S or Hipraviar-CLON	NCD	Eye drop /Spray or Drinking water
10 wks	Hiprapox	Fowl Pox	Wing stab



Murphy Chemicals E.A Ltd

Baba Dogo Road- Ruaraka P.O. Box 20495-00200 Nairobi, Kenya Tel:+254-20-2017992,
Cellphone: +254-722-570768 / 254-733-610860 Fax:+254-20-8562199
E-mail:murphy@murphychemicals.co.ke. Website: www.murphychemicals.co.ke



**NB: Vaccinate against NCD every 6-8 weeks with
Hipraviar-S or Hipraviar-CLON**

